

# Unlocking Neutron Stars as Probes of Fundamental Physics

## Tim Linden

Physics and Astronomy Colloquium  
University of Hawai'i at Mānoa  
February 7, 2019




THE OHIO STATE UNIVERSITY  
CENTER FOR COSMOLOGY AND  
ASTROPARTICLE PHYSICS



# Gravitational Dark Matter


---



- **Dark Matter is:**
  - **Dark**
  - **Cold**
  - **Stable**
  - **Prevalent**



# Gravitational Probes of Dark Matter






# Neutron Stars: The Big and the Small

---

- **Big:**  $\sim 1.4 M_{\odot}$
- **Small:** Compressed into 10 km
- **Big:** Can spin up to  $700 \text{ s}^{-1}$  (0.2 c at surface)
- **Small:** Oblate spheroid to  $< 1$  part in a million





# Neutron Stars: Precision Physics

- Neutron star spin among the best measured quantities in physics.

**PSR J1713+0747**

$$F = 218.8118437960826270 \pm 0.000000000000000988 \text{ s}^{-1}$$

$$F' = -4.083888637248 \pm 0.0000143324982645 \times 10^{-16} \text{ s s}^{-1}$$




# A Dipole Model

- Can precisely measure the magnetic field:

$$\frac{dE}{dt} \propto -\omega^4 R^6 B^2 \sin^2 \alpha$$
$$B \sim 3.3 \times 10^{19} \left[ P^2 \left( \frac{1}{P} \frac{dP}{dt} \right) \right]^{1/2} \text{ G}$$

- And approximately measure the age:


$$\tau = \frac{P}{2(dP/dt)}$$





# Curvature Radiation

- Changing magnetic field produces electric field




- 1000 PV potential available to accelerate particles








# Multiwavelength Emission





credit: Dave Thompson







# A Window Into Extreme Physics





# A Window Into General Relativity

THE ASTROPHYSICAL JOURNAL LETTERS, 848:L12 (59pp), 2017 October 20

© 2017. The American Astronomical Society. All rights reserved.

OPEN ACCESS

<https://doi.org/10.3847/2041-8213/aa91c9>



## Multi-messenger Observations of a Binary Neutron Star Merger

LIGO Scientific Collaboration and Virgo Collaboration, Fermi GBM, INTEGRAL, IceCube Collaboration, AstroSat Cadmium Zinc Telluride Imager Team, IPN Collaboration, The Insight-Hxmt Collaboration, ANTARES Collaboration, The Swift Collaboration, AGILE Team, The 1M2H Team, The Dark Energy Camera GW-EM Collaboration and the DES Collaboration, The DLT40 Collaboration, GRAVITA: GRAvitational Wave Inaf TeAm, The Fermi Large Area Telescope Collaboration, ATCA: Australia Telescope Compact Array, ASKAP: Australian SKA Pathfinder, Las Cumbres Observatory Group, OzGrav, DWF (Deeper, Wider, Faster Program), AST3, and CAASTRO Collaborations, The VINROUGE Collaboration, MASTER Collaboration, J-GEM, GROWTH, JAGWAR, Caltech-NRAO, TTU-NRAO, and NuSTAR Collaborations, Pan-STARRS, The MAXI Team, TZAC Consortium, KU Collaboration, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS: Transient Robotic Observatory of the South Collaboration, The BOOTES Collaboration, MWA: Murchison Widefield Array, The CALET Collaboration, IKI-GW Follow-up Collaboration, H.E.S.S. Collaboration, LOFAR Collaboration, LWA: Long Wavelength Array, HAWC Collaboration, The Pierre Auger Collaboration, ALMA Collaboration, Euro VLBI Team, Pi of the Sky Collaboration, The Chandra Team at McGill University, DFN: Desert Fireball Network, ATLAS, High Time Resolution Universe Survey, RIMAS and RATIR, and SKA South Africa/MeerKAT  
(See the end matter for the full list of authors.)

Received 2017 October 3; revised 2017 October 6; accepted 2017 October 6; published 2017 October 16

### Abstract


On 2017 August 17 a binary neutron star coalescence candidate (later designated GW170817) with merger time 12:41:04 UTC was observed through gravitational waves by the Advanced LIGO and Advanced Virgo detectors. The *Fermi* Gamma-ray Burst Monitor independently detected a gamma-ray burst (GRB 170817A) with a time delay of  $\sim 1.7$  s with respect to the merger time. From the gravitational-wave signal, the source was initially localized to a sky region of  $31 \text{ deg}^2$  at a luminosity distance of  $40^{+8}_{-8}$  Mpc and with component masses consistent with neutron stars. The component masses were later measured to be in the range  $0.86$  to  $2.26 M_{\odot}$ . An extensive observing campaign was launched across the electromagnetic spectrum leading to the discovery of a bright optical transient (SSS17a, now with the IAU identification of AT 2017gfo) in NGC 4993 (at  $\sim 40$  Mpc) less than 11 hours after the merger by the One-Meter, Two Hemisphere (1M2H) team using the 1 m Swope Telescope. The optical transient was independently detected by multiple teams within an hour. Subsequent observations targeted the object and its environment. Early ultraviolet observations revealed a blue transient that faded within 48 hours. Optical and infrared observations showed a redward evolution over  $\sim 10$  days. Following early non-detections, X-ray and radio emission were discovered at the transient's position  $\sim 9$  and  $\sim 16$  days, respectively, after the merger. Both the X-ray and radio emission likely arise from a physical process that is distinct from the one that generates the UV/optical/near-infrared emission. No ultra-high-energy gamma-rays and no neutrino candidates consistent with the source were found in follow-up searches. These observations support the hypothesis that GW170817 was produced by the merger of two neutron stars in NGC 4993 followed by a short gamma-ray burst (GRB 170817A) and a kilonova/macronova powered by the radioactive decay of *r*-process nuclei synthesized in the ejecta.

**Key words:** — gravitational waves — neutron stars — kilonova


gravitational waves of *r*-process nuclei synthesized in the ejecta. NGC 4993 followed by a short gamma-ray burst (GRB 170817A) and a kilonova/macronova powered by the radioactive decay of *r*-process nuclei synthesized in the ejecta. These observations support the hypothesis that GW170817 was produced by the merger of two neutron stars in NGC 4993 followed by a short gamma-ray burst (GRB 170817A) and a kilonova/macronova powered by the radioactive decay of *r*-process nuclei synthesized in the ejecta.




# A Window Into Astrophysics



Massive Stars



Shock Acceleration



Positron Excess



# A Window Into Fundamental Physics


---

- **Sensitive probes of rare processes:**
  1. **Nuclear densities over macroscopic distances**
  2. **Strongest magnetic fields in the universe**
- **Precise measurements are possible**



# The Program

---






# A Window Into Fundamental Physics

---

- Sensitive probes of rare processes:
  1. Nuclear densities over macroscopic distances
  2. Strongest magnetic fields in the universe
- Precise measurements are possible




# Searching for Dark Matter Interactions






# Direct Detection: Experimental Efforts









# Neutron Stars: The Optimal Direct Detection Experiment



## LUX


- 100 kg
- 1000 days

**$10^5$  kg day**

## Neutron Star

- $10^{30}$  kg
- $10^{12}$  days


**$10^{42}$  kg day**





# Neutron Stars: The Optimal Direct Detection Experiment

- Neutron stars are so dense that they are optically thick to dark matter




$$\sigma_{\text{sat}}^{\text{single}} \simeq \pi R^2 m_n / M \simeq 2 \times 10^{-45} \text{ cm}^2 \left( \frac{1.5 M_{\odot}}{M} \right) \left( \frac{R}{10 \text{ km}} \right)^2$$

- This saturates the sensitivity of neutron stars to dark matter



# Neutron Stars: Astrophysics Enhancements

- Neutron stars gravitationally attract nearby dark matter



Capture radius is approximately  $1 R_0$

$$b_{\max} = \left( \frac{2GM R}{v_x^2} \right)^{1/2} \left( 1 - \frac{2GM}{R} \right)^{-1/2}$$


$$\dot{m} = \pi b_{\max}^2 v_x \rho_x,$$

- Interaction scales as  $v_x^{-1}$ , very sensitive to slowly moving dark matter



# Neutron Stars: Astrophysics Enhancements

- Neutron stars are a dark matter collider



When dark matter hits the neutron star surface it is moving relativistically:

$$v_{esc} = \sqrt{\frac{2GM}{r}} \sim 0.7c$$


- Can probe p-wave suppressed dark matter or dark matter mass splittings



# Neutron Stars: Particle Physics Complications

Typical NS neutron momentum is:

$$p_{F,n} \simeq 0.45 \text{ GeV} \left( \rho_{NS} / (4 \times 10^{38} \text{ GeV cm}^{-3}) \right)$$



This suppresses the interaction cross-section for low mass DM:

$$\sigma_{\text{sat}}^{\text{Pauli}} \simeq \pi R^2 m_n p_f / (M \gamma m_x v_{\text{esc}}) \simeq 2 \times 10^{-45} \text{ cm}^2 \left( \frac{\text{GeV}}{m_x} \right) \left( \frac{1.5 M_\odot}{M} \right) \left( \frac{R}{10 \text{ km}} \right)^2.$$



# Neutron Stars: Particle Physics Complications

Dark Matter kinetic energy lost in a scatter with a proton is:

$$E_{loss} = \frac{2m_p}{m_\chi} (m_\chi v_\chi^2)$$




Very heavy dark matter requires multiple interactions:

$$\sigma_{\text{sat}}^{\text{multi}} \simeq 2 \times 10^{-45} \text{ cm}^2 \left( \frac{m_\chi}{\text{PeV}} \right) \left( \frac{1.5 \text{ M}_\odot}{M} \right) \left( \frac{R}{10 \text{ km}} \right)^2.$$



# Neutron Stars: Particle Physics Complications







# **Detecting Dark Matter Scattering in Neutron Stars**

## **Part I: Neutron Star Heating**



# Dark Matter Induced Heating

Energy In = Energy Out



**DM-NS collisions impart significant energy into the NS:**

$$E_s \simeq m_x (\gamma - 1)$$



**This induces blackbody emission of luminosity:**

$$\dot{E}_k = \frac{E_s \dot{m}}{m_x} f \simeq 1.4 \times 10^{25} \text{ GeV s}^{-1} \left( \frac{f}{1} \right),$$



# Dark Matter Induced Heating

Energy In = Energy Out



If Dark Matter subsequently annihilates, additional energy is injected (de Lavellez & Fairbairn (1004.0629))



$$E_s \simeq m_x (\gamma - 1)$$




# Detecting Hot Neutron Stars

- Thermal emission detected from young neutron stars
- Older neutron stars continue cooling
- Dark matter sets a minimum temperature of  $\sim 2000$  K ( $10^{22}$  erg)

Potekhin & Chabrier (1711.07662)











# Detecting Thermal Emission

- Observations at 2000 K require infrared telescopes



**JWST**  
**10 nJy in  $10^4$  s**



**GMT**  
**0.5 nJy in  $10^5$  s**

- A pulsar at 10 pc would have a flux of  $\sim 2$  nJy at 2 microns



# What Do We Need?

---

1. A nearby pulsar (10-20 pc).
  - Closest observed pulsar: 90 pc (PSR B1055-52)
  - Average Distance to nearest NS: 10 pc (Sartore et al. 0908.3182)
2. A model to separate thermal from pulsed emission
3. Constraints on thermal injection sources, e.g. gas accretion and magnetic heating.



# **Detecting Dark Matter Scattering in Neutron Stars**

## **Part II: Dark Matter Collapse**



# The Secret Life of Dark Matter Inside a Neutron Star

---

- **Capture** - DM hits neutron and elastically scatters
- **Thermalization** - Trapped dark matter thermalizes with neutron superfluid. If dark matter can annihilate, it will.
- **Collapse** - Dark matter degeneracy pressure not capable of preventing collapse.

Bramante & TL (1405.1031)

Bramante & TL (1601.06784)

Bramante, TL, Tsai (1706.00001)







# Dark Matter Parameter Space

---


- Requires dark matter to be non-annihilating.

- **PeV Fermionic Dark Matter**

$$M_{crit}^{ferm} \simeq M_{pl}^3 / m_X^2$$

- **Bosonic Dark Matter**
- **Attractive Self-Interacting Dark Matter**









# Strong Constraints are Possible

McDermott et al. (1103.5472)





# A Signal

**10% of Star Formation in central 200 pc of Milky Way**

**Only one (very young) pulsar detected**

**The Missing Pulsar problem!**

**Massive Star Formation  
in the Galactic Center**  
*By Don F. Figer*  
Rochester Institute of Technology, Rochester, NY

The Galactic center is a hotbed of star formation activity, containing more massive young stars with initial masses up to 100 solar masses than anywhere else in the Galaxy. This review concerns the young stellar population in the region surrounding the central black hole, and the bulk of the massive stellar clusters, the population of younger stars in the stars surrounding the central black hole, and the fossil record in the Galactic center suggests that the recently formed present-day examples of similar populations that must have been formed in episodes stretching back to the time period when the Galaxy was forming.

**THE PECULIAR PULSAR POPULATION OF THE CENTRAL PARSEC**  
*JASON DEXTER*  
Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

**RYAN M. O'LEARY**  
Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA  
*Draft version April 14, 2018*

**ABSTRACT**  
The Galactic center black hole, Sgr A\*, would be potential probes of its environment. It can be used to test general relativity. Despite predictions of thousands of millisecond pulsars in the Galactic center, none have been discovered. The discovery of radio pulsations from a highly magnetized neutron star in the temporal pulsations from a highly magnetized neutron star in the Galactic center has been that hyperstrong temporal pulsations from a highly magnetized neutron star in the Galactic center is much weaker than ordinary pulsars. The discovery of radio pulsations from a highly magnetized neutron star in the Galactic center is much weaker than ordinary pulsars. The discovery of radio pulsations from a highly magnetized neutron star in the Galactic center is much weaker than ordinary pulsars.

**Introduction**  
The Galactic center (GC) is an important region for understanding the formation and evolution of galaxies. It contains a supermassive black hole, surrounded by a dense concentration of stars and gas. The GC is also a site of intense star formation, producing many massive stars and supernovae. In this paper, we discuss the properties of the GC and the challenges associated with studying it.

# 10% of Star Formation in central 200 pc of Milky Way

## Only one (very young) pulsar detected

# The Missing Pulsar problem!

# **Massive Star Formation in the Galactic Center**

By Don F. Figer  
Institute of Technology

# Galactic Cent

By Don F. Figer

Rochester Institute of Technology, Rochester, NY

is a hotbed of star formation acti  
ree of the most massive star  
contains more stars  
review con  
star f

The Galactic center is a hotbed of star formation activity, containing more massive young stars with initial masses up to 100 solar masses than anywhere else in the Galaxy. This review concerns the population of younger stars in the central black hole, and the bulk of the massive stellar clusters surrounding it. The fossil record in the Galactic center suggests that the recently formed stars are part of a long-term process present-day examples of similar populations that must have been forming since the time period when the Galaxy was first assembled.

**Introduction**

The Galactic center (GC) is an important astrophysical environment. It contains a tiny fraction of the total mass of the Galaxy, but it is a rich source of information about the processes that shape the evolution of galaxies.

roduction  
Galact:

the Galactic center (GC) is an extremely dense region of the Galaxy where billions of stars and a supermassive black hole (SMBH) reside. It is the heart of the Galaxy, where the gravitational pull is so strong that it holds the entire Galaxy together. The GC is also the site of many of the most energetic and violent events in the Galaxy, including supernovae, gamma-ray bursts, and the formation of new stars. The discovery of the first pulsar in the GC, PSR B1509-58, in 1991, was a major breakthrough in our understanding of the inner workings of the Galaxy. This pulsar, located about 26,000 light-years from Earth, is the only one of its kind in the GC and is thought to be a neutron star, the collapsed core of a massive star that has exploded in a supernova. The discovery of PSR B1509-58 opened up a new window into the study of the GC and its inhabitants, and it continues to be a subject of intense research and discovery.

THE PECULIAR PULSAR POPULATION OF THE CENTRAL PARSEC

JASON DEXTER  
Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

RYAN M. O'LEARY  
University of California, Berkeley, CA 94720-3411

Draft version April 14, 2018

ABSTRACT

\* would be potential p...  
... Despite predi...  
... none h...  
... strong

POPULATION  
JASON DEXTER  
City of California

RYAN M. O'LEARY  
University of California  
April 11

RYAN M. O'LEARY  
University of California, Berkeley  
Draft version April 14, 2018

**ABSTRACT**

A\*, working on the

ABSTRACT

**R PULSAR POPULATION**

JASON D. MILLER  
Department of Astronomy, University of California, Berkeley, CA

RYAN M. O'LEARY  
Department of Astronomy, University of California, Berkeley, CA  
*Draft version April 14, 2018*

**ABSTRACT**

The Galactic center black hole, Sgr A\*, would be potential probes of it  
can be used to test general relativity. Despite predictions of  
millisecond pulsars in the Galactic center, none have been  
discovered. The discovery of radio pulsations from a highly ma  
the temporal scattering is much weaker  
ary pulsar population is the most  
"magnetar" in the Galactic center.  
ordinary pulsar  
could be c  
mass



# A Signal

10% of Star Formation in central 200 pc of Milky Way

Only one (very young) pulsar detected

## The Missing Pulsar problem!

**ALERT: THIS BIKE IS STOLEN**



UP32 CW 5418

This N.O.W. bikers group bike has been stolen from Gautam Nagar area in New Delhi on 10th December 2010. Any one who sees or have any info about this bike, please contact Delhi Police on 100 or the vehicle owner Aditya Ratti at 9582126004. Many Thanks

E CENTRAL PARSEC  
Berkeley, CA 94720-3411, USA

14, 2018

ACT  
A\*, would be potential probes of it  
relativity. Despite predictions of  
Galactic center, none have been  
as been that hyperstrong temporal  
radio pulsations from a highly ma  
temporal scattering is much weak  
ary pulsar population is the most  
"magnetar formation in  
ordinary pulsar  
could be c  
mass

Roche  
The Galactic center is a ho  
formation site and three of t  
a rich environment, it contains  
else in the Galaxy. This review  
as it relates to massive star form  
massive stellar clusters, the popul  
the stars surrounding the central b  
The fossil record in the Galactic cent  
present-day examples of similar popula  
isodes stretching back to the time pe  
Introduction  
Galactic center (GC) is an  
tion models. It con  
ly a tiny f



# The Missing Pulsar Problem

- **High Dark Matter density near the GC.**


$$\rho(r) = \frac{\rho_0}{\frac{r}{R_s} \left( 1 + \frac{r}{R_s} \right)^2}$$

- **GC NS collapse in  $\sim 10^5$  yr while nearby NS remain.**


- **Constrains cross-section to within a few orders of magnitude.**













---


**Hard to detect dark matter through a null observation.**

**What observations stand as evidence of dark matter  
induced neutron star collapse?**



# An Electromagnetic Signal

---






# An Electromagnetic Signal

- **Gamma-Ray Bursts (observed by Fermi)**
- **Optical emission from the decay of r-process elements**
- **Fast Radio Bursts are potentially correlated with NS mergers.**


## Fermi GBM Collaboration (1710.05834)






# An Electromagnetic Signal

No DM Induced Collapse




DM Induced Collapse






# An Electromagnetic Signal

## No DM Induced Collapse




## DM Induced Collapse






# New Phenomena



## Merger Kilonovae

Electromagnetic signals  
and gravitational waves  
jointly identified.


(proportional to  $\rho_{\text{DM}}^{-1}$ )



## Quiet Kilonovae

Electromagnetic signals  
without gravitational  
waves.

(proportional to  $\rho_{\text{DM}}$ ).



## Dark Mergers

Gravitational waves  
without any  
electromagnetic signal.

(proportional to  $\rho_{\text{DM}}$ ).




# Two Methods

---

- **Two methods to isolate dark matter signal:**
  - 1. Look in regions where dark matter induced signal is dominant (e.g. dwarf galaxies)**
  - 2. Examine the spatial morphology of events in and extract dark matter density profile.**




# Merger Kilonovae






# Constraining Dark Matter - Merger Kilonovae






# Fast Radio Bursts or Quiet Kilonovae





# Finding Dark Matter - Fast Radio Bursts





# What Do We Need?

---

- 1. New Observations of NS Mergers (gravitational waves, electromagnetic emission, fast radio bursts).**
- 2. Localization of the electromagnetic signatures within galaxies.**
- 3. Improved models for the electromagnetic signals from dark matter induced NS collapse.**



# A Window Into Fundamental Physics

---


- **Sensitive probes of rare processes:**
  1. Nuclear densities over macroscopic distances.
  2. **Strongest magnetic fields in the universe.**
- **Precise measurements are possible.**



# Detecting Axion Dark Matter

- We can search for the resonant decay to photons:

$$P_{\text{SIG}} = \eta g_{a\gamma\gamma}^2 \left( \frac{\rho_a}{m_a} \right) B_0^2 V C Q_L$$





# Neutron Stars: The Optimal Axion Laboratory

$$P_{a\gamma} \sim g_{a\gamma\gamma}^2 \mathbf{B}^2 L^2$$



## ADMX


- 10 T
- 1 m<sup>2</sup>

100 T<sup>2</sup> m<sup>2</sup>

## Neutron Star

- 10<sup>10</sup> T
- 10<sup>8</sup> m<sup>2</sup>

10<sup>28</sup> T<sup>2</sup> m<sup>2</sup>






# Neutron Stars: The Optimal Axion Laboratory

- Resonant interactions occur when plasma frequency equals the axion mass:

$$\omega_p \approx (1.5 \times 10^2 \text{ GHz}) \sqrt{\left(\frac{B_z}{10^{14} \text{ G}}\right) \left(\frac{1 \text{ sec}}{P}\right)}$$

$$= 6 \times 10^{-4} \text{ eV}$$

Mitra et al. (1510.00103)




- Need detailed model of NS magnetic fields.

$$r_c(\theta, \theta_m, t) = 224 \text{ km} \times |3 \cos \theta \hat{\mathbf{m}} \cdot \hat{\mathbf{r}} - \cos \theta_m|^{1/3} \times \left(\frac{r_0}{10 \text{ km}}\right) \times \left[\frac{B_0}{10^{14} \text{ G}} \frac{1 \text{ sec}}{P} \left(\frac{1 \text{ GHz}}{m_a}\right)^2\right]^{1/3}.$$

Hook et al. (1804.03145)



# Neutron Stars: The Optimal Axion Laboratory



- Can place complementary constraints on the QCD axion.
- Specific to models where axions are the dark matter.



# What Do We Need?


---

- 1. Nearby, highly-magnetized pulsar.**
- 2. Better models of the pulsar magnetic field.**
- 3. Sensitive observations of radio lines (different techniques than traditional pulsar searches).**




# Finding the Right Neutron Star

---






# Radio Pulses: A Blessing and a Curse




Harding (2016; J Plasma Phys 82)






# A New Method for Detecting Invisible Pulsars

---





# A New Method for Detecting Invisible Pulsars



2° ~ 10 pc

Geminga

PSR B0656+14



# Astrophysical Implications of TeV Halos


---

- **TeV halo observations solve many astrophysical puzzles**
- **Prove that pulsars produce the positron excess**  
(Hooper, Cholis, TL, Fang 1702.08436)
- **Explain the TeV gamma-ray excess**  
(TL & Buckman 1707.01905)
- **Explain inhomogeneities in cosmic-ray diffusion ,**  
(Hooper & TL 1711.07482) (Evoli, TL, Morlino, TBS)
- **Explain TeV gamma-rays from the Galactic center**  
(Hooper et al. 1705.09293)




# Discovering Pulsars at TeV Energies

- Tauris and Manchester (1998) calculated the beaming angle from a population of young and middle-aged pulsars.
- This varies between 15-30%.
- $1/f$  pulsars are unseen in radio surveys.





# Discovering Invisible Pulsars at TeV Energies



## The H.E.S.S. Galactic plane survey


H.E.S.S. Collaboration, H. Abdalla<sup>1</sup>, A. Abramowski<sup>2</sup>, F. Aharonian<sup>3,4,5</sup>, F. Ait Benkhali<sup>3</sup>, E.O. Angüner<sup>21</sup>, M. Arakawa<sup>43</sup>, M. Arrieta<sup>15</sup>, P. Aubert<sup>24</sup>, M. Backes<sup>8</sup>, A. Balzer<sup>9</sup>, M. Barnard<sup>1</sup>, Y. Becherini<sup>10</sup>, J. Becker Tjus<sup>11</sup>, D. Berge<sup>12</sup>, S. Bernhard<sup>13</sup>, K. Bernlöhr<sup>3</sup>, R. Blackwell<sup>14</sup>, M. Böttcher<sup>1</sup>, C. Boisson<sup>15</sup>, J. Bolmont<sup>16</sup>, S. Bonnefoy<sup>37</sup>, P. Bortas<sup>3</sup>, J. Bregeon<sup>17</sup>, F. Brun<sup>18</sup>, P. Brun<sup>18</sup>, M. Bryan<sup>9</sup>, M. Büchele<sup>36</sup>, T. Bulik<sup>19</sup>, M. Capasso<sup>29</sup>, S. Carrigan<sup>3,48</sup>, S. Caroff<sup>30</sup>, A. Carosi<sup>24</sup>, S. Casanova<sup>21,3</sup>, M. Cerruti<sup>16</sup>, N. Chakraborty<sup>3</sup>, R.C.G. Chaves<sup>17,22</sup>, A. Chen<sup>23</sup>, J. Chevalier<sup>24</sup>, S. Colafrancesco<sup>23</sup>, B. Condon<sup>26</sup>, J. Conrad<sup>27,28</sup>, I.D. Davids<sup>8</sup>, J. Decock<sup>18</sup>, C. Deil<sup>3</sup>, J. Devin<sup>17</sup>, P. deWilt<sup>14</sup>, L. Dirson<sup>2</sup>, A. Djannati-Atai<sup>31</sup>, W. Domainko<sup>3</sup>, A. Donath<sup>3</sup>, L.O'C. Drury<sup>4</sup>, K. Dutson<sup>33</sup>, J. Dyks<sup>34</sup>, T. Edwards<sup>3</sup>, K. Egberts<sup>35</sup>, P. Eger<sup>3</sup>, G. Emery<sup>16</sup>, J.-P. Ernenwein<sup>20</sup>, S. Eschbach<sup>36</sup>, C. Farnier<sup>27,10</sup>, S. Fegan<sup>30</sup>, M.V. Fernandes<sup>2</sup>, A. Fiasson<sup>24</sup>, G. Fontaine<sup>30</sup>, A. Förster<sup>3</sup>, S. Funk<sup>36</sup>, M. Füßling<sup>37</sup>, S. Gabici<sup>31</sup>, Y.A. Gallant<sup>17</sup>, T. Garrigoux<sup>1</sup>, H. Gast<sup>3,49</sup>, F. Gaté<sup>24</sup>, G. Giavitto<sup>37</sup>, B. Giebels<sup>30</sup>, D. Glawion<sup>25</sup>, J.F. Glicenstein<sup>18</sup>, D. Gottschall<sup>29</sup>, M.-H. Grondin<sup>26</sup>, J. Hahn<sup>3</sup>, M. Haupt<sup>37</sup>, J. Hawkes<sup>14</sup>, G. Heinzlmann<sup>2</sup>, G. Henri<sup>32</sup>, G. Hermann<sup>3</sup>, J.A. Hinton<sup>3</sup>, W. Hofmann<sup>3</sup>, C. Hoischen<sup>35</sup>, T. L. Holch<sup>7</sup>, M. Holler<sup>13</sup>, D. Horns<sup>2</sup>, A. Ivascenko<sup>1</sup>, H. Iwasaki<sup>43</sup>, A. Jacholkowska<sup>16</sup>, M. Jamroz<sup>38</sup>, D. Jankowsky<sup>36</sup>, F. Jankowsky<sup>25</sup>, M. Jingo<sup>23</sup>, L. Jouvin<sup>31</sup>, I. Jung-Richardt<sup>36</sup>, M.A. Kastendieck<sup>2</sup>, K. Katarzynski<sup>39</sup>, M. Katsuragawa<sup>44</sup>, U. Katz<sup>36</sup>, D. Kerszberg<sup>16</sup>, D. Khangulyan<sup>43</sup>, B. Khélifi<sup>31</sup>, J. King<sup>3</sup>, S. Klepser<sup>37</sup>, D. Klockov<sup>29</sup>, W. Kluźniak<sup>34</sup>, Nu. Komin<sup>23</sup>, K. Kosack<sup>18</sup>, S. Krakau<sup>11</sup>, M. Kraus<sup>36</sup>, P.P. Krüger<sup>1</sup>, H. Laffon<sup>3</sup>, G. Lamanna<sup>24</sup>, J. Lau<sup>14</sup>, J.-P. Lees<sup>24</sup>, J. Lefaucheur<sup>15</sup>, A. Lemièr<sup>31</sup>, M. Lemoine-Goumard<sup>26</sup>, J.-P. Lenain<sup>16</sup>, E. Leser<sup>35</sup>, T. Lohse<sup>7</sup>, M. Lorentz<sup>18</sup>, R. Liu<sup>3</sup>, R. López-Coto<sup>3</sup>, I. Lypova<sup>37</sup>, V. Marandon<sup>3</sup>, D. Malyshev<sup>29</sup>, A. Marcowith<sup>17</sup>, C. Mariaud<sup>30</sup>, R. Marx<sup>3</sup>, G. Maurin<sup>24</sup>, N. Maxted<sup>14,45</sup>, M. Mayer<sup>7</sup>, P.J. Meintjes<sup>40</sup>, M. Meyer<sup>27</sup>, A.M.W. Mitchell<sup>3</sup>, R. Moderski<sup>34</sup>, M. Mohamed<sup>25</sup>, L. Mohrmann<sup>36</sup>, K. Morá<sup>27</sup>, E. Moulin<sup>18</sup>, T. Murach<sup>37</sup>, S. Nakashima<sup>44</sup>, M. de Naurois<sup>30</sup>, H. Ndiyavala<sup>1</sup>, F. Niederwanger<sup>13</sup>, J. Niemiec<sup>21</sup>, L. Oakes<sup>7</sup>, P. O'Brien<sup>33</sup>, H. Odaka<sup>44</sup>, S. Ohm<sup>37</sup>, M. Ostrowski<sup>38</sup>, I. Oya<sup>37</sup>, M. Padovani<sup>17</sup>, M. Panter<sup>3</sup>, R.D. Parsons<sup>3</sup>, M. Paz Arribas<sup>7</sup>, N.W. Pekeur<sup>1</sup>, G. Pelletier<sup>32</sup>, C. Perennes<sup>16</sup>, P.-O. Petrucci<sup>32</sup>, B. Peyaud<sup>18</sup>, Q. Piel<sup>24</sup>, S. Pita<sup>31</sup>, V. Poireau<sup>24</sup>, H. Poon<sup>3</sup>, D. Prokhorov<sup>10</sup>, H. Prokoph<sup>12</sup>, G. Pühlhofer<sup>29</sup>, M. Punch<sup>31,10</sup>, A. Quirrenbach<sup>25</sup>, S. Raab<sup>36</sup>, R. Rauth<sup>13</sup>, A. Reimer<sup>13</sup>, O. Reimer<sup>13</sup>, M. Renaud<sup>17</sup>, R. de los Reyes<sup>3</sup>, F. Rieger<sup>3,41</sup>, L. Rinchiuso<sup>18</sup>, C. Romoli<sup>4</sup>, G. Rowell<sup>14</sup>, B. Rudak<sup>34</sup>, C.B. Rulten<sup>15</sup>, S. Safi-Harb<sup>50</sup>, V. Sahakian<sup>6,5</sup>, S. Saito<sup>43</sup>, D.A. Sanchez<sup>24</sup>, A. Santangelo<sup>29</sup>, M. Sasaki<sup>36</sup>, M. Schandri<sup>36</sup>, R. Schlickeiser<sup>11</sup>, F. Schlüssler<sup>18</sup>, A. Schulz<sup>37</sup>, U. Schwanke<sup>7</sup>, S. Schwemmer<sup>25</sup>, M. Seglar-Arroyo<sup>18</sup>, M. Settimo<sup>16</sup>, A.S. Seyffert<sup>1</sup>, N. Shafi<sup>23</sup>, I. Shilon<sup>36</sup>, K. Shiningayamwe<sup>8</sup>, R. Simoni<sup>9</sup>, H. Sol<sup>15</sup>, F. Spanier<sup>1</sup>, M. Spir-Jacob<sup>31</sup>, L. Stawarz<sup>38</sup>, R. Steenkamp<sup>8</sup>, C. Stegmann<sup>35,37</sup>, C. Steppa<sup>35</sup>, I. Sushch<sup>1</sup>, T. Takahashi<sup>44</sup>, J.-P. Tavernier<sup>16</sup>, T. Tavernier<sup>31</sup>, A.M. Taylor<sup>37</sup>, R. Terrier<sup>31</sup>, L. Tibaldo<sup>3</sup>, D. Tiziani<sup>36</sup>, M. Tluczykont<sup>2</sup>, C. Trichard<sup>20</sup>, M. Tsiros<sup>17</sup>, N. Tsuji<sup>43</sup>, R. Tufts<sup>3</sup>, Y. Uchiyama<sup>43</sup>, D.J. van der Walt<sup>1</sup>, C. van Eldik<sup>36</sup>, C. van Rensburg<sup>1</sup>, B. van Soelen<sup>40</sup>, G. Vasileiadis<sup>17</sup>, J. Veh<sup>36</sup>, C. Venter<sup>1</sup>, A. Viana<sup>3,46</sup>, P. Vincent<sup>16</sup>, J. Vink<sup>9</sup>, F. Voisin<sup>14</sup>, H.J. Völk<sup>3</sup>, Z. Wadiasingh<sup>1</sup>, S.J. Wagner<sup>25</sup>, P. Wagner<sup>7</sup>, R.M. Wagner<sup>27</sup>, R. White<sup>3</sup>, A. Wierchowska<sup>21</sup>, P. Willmann<sup>36</sup>, A. Wörnlein<sup>36</sup>, D. Wouters<sup>18</sup>, R. Yang<sup>3</sup>, D. Zaborov<sup>30</sup>, M. Zacharias<sup>1</sup>, R. Zanin<sup>3</sup>, A.A. Zdziarski<sup>34</sup>, A. Zech<sup>15</sup>, F. Zefi<sup>30</sup>, A. Ziegler<sup>36</sup>, J. Zorn<sup>3</sup>, and N. Zywucka<sup>38</sup>

(Affiliations can be found after the references)

April 10, 2018


### ABSTRACT

We present the results of the most comprehensive survey of the Galactic plane in very high-energy (VHE)  $\gamma$ -rays, including a public release of Galactic sky maps, a catalog of VHE sources, and the discovery of 16 new sources of VHE  $\gamma$ -rays. The High Energy Spectroscopic System (H.E.S.S.) Galactic plane survey (HGPS) was a decade-long observation program carried out by the H.E.S.S. I array of Cherenkov telescopes in Namibia from 2004 to 2013. The observations amount to nearly 2700 h of quality-selected data, covering the Galactic plane at longitudes from  $\ell = 250^\circ$  to  $65^\circ$  and latitudes  $|b| \leq 3^\circ$ . In addition to the unprecedented spatial coverage, the HGPS also features a relatively high angular resolution ( $0.08^\circ \approx 5$  arcmin mean point spread function 68% containment radius), sensitivity ( $\lesssim 1.5\%$  Crab flux for point-like sources), and energy range (0.2 to 100 TeV). We constructed a catalog of VHE  $\gamma$ -ray sources from the HGPS data set with a systematic procedure for both source detection and characterization of morphology and spectrum. We present this likelihood-based method in detail, including the introduction of a model component to account for unresolved, large-scale emission along the Galactic plane. In total, the resulting HGPS catalog contains 78 VHE sources, of which 14 are not reanalyzed here, for example, due to their complex morphology, namely shell-like sources and the Galactic center region. Where possible, we provide a firm identification of the VHE source or plausible associations with sources in other astronomical catalogs. We also studied



02432v1 [astro-ph.HE] 6 Apr 2018







# What Do We Need?

---

- 1. Continued observations of TeV halos.**
- 2. A model for the confinement and emission of electrons in TeV halos.**
- 3. A method for precisely determining the pulsar position within the TeV halo.**




# A Model for TeV Halos

Evoli, TL, Morlino (1807.09263)

- Early results indicate that **pulsars themselves can confine electrons to produce TeV halo emission.**

- Analog with cosmic-ray confinement in supernova remnants.



- More detailed models including reacceleration and joint supernova/pulsar emission are necessary.




# The Program

---

## 1. Understand Dark Matter/Neutron Star Interactions

- Can already set strong constraints on some models
  1. Asymmetric Dark Matter
  2. Axions
- Can probe extremely generic dark matter models.






# The Program

---

## 1. Understand Dark Matter/Neutron Star Interactions

- Can already set strong constraints on some models
  1. Asymmetric Dark Matter
  2. Axions
- Can probe extremely generic dark matter models.




## 2. Differentiate dim dark matter signals from astrophysics

- Need detailed models of neutron star physics.
- Requires observations of pulsars with “special” attributes
  1. Nearby
  2. Strong Magnetic Fields
  3. Not Beamed Towards Earth




















# Conclusions

---

- **Pulsars have unique characteristics that are optimally suited for new physics searches.**
- **Early studies can set strong constraints on the asymmetric dark matter and axion parameter spaces.**
- **Our observational techniques are in their infancy. The next decade will revolutionize the field in several directions.**




**Extra Slides**



# Particle Physics Mash-Up

M A S H U P

- Can also use neutron star cooling curves to place limits on the axion cross-section.
- Observations of the Cassiopeia A NS, with a known age of 337 years, rule out  $f_a < 5 \times 10^8$  GeV.



Hamaguchi et al. (1806.07151)



# Discovering Pulsars at TeV Energies

- 5 / 39 sources in the 2HWC catalog are correlated with bright, middle-aged (100 — 400 kyr) pulsars.


2HWC Name	ATNF Name	Distance (kpc)	Angular Separation	Projected Separation	Expected Flux ( $\times 10^{-15}$ )	Actual Flux ( $\times 10^{-15}$ )	Flux Ratio	Expected Extension	Actual Extension	Age (kyr)	Chance Overlap
J0700+143	B0656+14	0.29	0.18°	0.91 pc	43.0	23.0	1.87	2.0°	1.73°	111	0.0
J0631+169	J0633+1746	0.25	0.89°	3.88 pc	48.7	48.7	1.0	2.0°	2.0°	342	0.0
J1912+099	J1913+1011	4.61	0.34°	27.36 pc	13.0	36.6	0.36	0.11°	0.7°	169	0.30
J2031+415	J2032+4127	1.70	0.11°	3.26 pc	5.59	61.6	0.091	0.29°	0.7°	181	0.002
J1831-098	J1831-0952	3.68	0.04°	2.57 pc	7.70	95.8	0.080	0.14°	0.9°	128	0.006

- 12 others with young pulsars
  - 2.3 chance overlaps
  - TeV emission may be contaminated by SNR

2HWC Name	ATNF Name	Distance (kpc)	Angular Separation	Projected Separation	Expected Flux ( $\times 10^{-15}$ )	Actual Flux ( $\times 10^{-15}$ )	Flux Ratio	Expected Extension	Actual Extension	Age (kyr)	Chance Overlap
J1930+188	J1930+1852	7.0	0.03°	3.67 pc	23.2	9.8	2.37	0.07°	0.0°	2.89	0.002
J1814-173	J1813-1749	4.7	0.54°	44.30 pc	243	152	1.60	0.11°	1.0°	5.6	0.61
J2019+367	J2021+3651	1.8	0.27°	8.48 pc	99.8	58.2	1.71	0.28°	0.7°	17.2	0.04
J1928+177	J1928+1746	4.34	0.03°	2.27 pc	8.08	10.0	0.81	0.11°	0.0°	82.6	0.002
J1908+063	J1907+0602	2.58	0.36°	16.21 pc	40.0	85.0	0.47	0.2°	0.8°	19.5	0.26
J2020+403	J2021+4026	2.15	0.18°	6.75 pc	2.48	18.5	0.134	0.23°	0.0°	77	0.01
J1857+027	J1856+0245	6.32	0.12°	13.24 pc	11.0	97.0	0.11	0.08°	0.9°	20.6	0.06
J1825-134	J1826-1334	3.61	0.20°	12.66 pc	20.5	249	0.082	0.14°	0.9°	21.4	0.14
J1837-065	J1838-0655	6.60	0.38°	43.77 pc	12.0	341	0.035	0.08°	2.0°	22.7	0.48
J1837-065	J1837-0604	4.78	0.50°	41.71 pc	8.3	341	0.024	0.10°	2.0°	33.8	0.68
J2006+341	J2004+3429	10.8	0.42°	80.07 pc	0.48	24.5	0.019	0.04°	0.9°	18.5	0.08



# Discovering Pulsars at TeV Energies




- Correcting for the beaming fraction implies that  $56^{+15}_{-11}$  TeV halos are currently observed by HAWC.
- However, only 39 HAWC sources total.
- Chance overlaps, SNR contamination must be taken into account.



# Discovering Pulsars at TeV Energies

- **Tentative Evidence that MSPs also produce these TeV halos.**
- **MSPs are the coldest and oldest pulsars – important for DM heating.**
- **Models indicate a MSP should exist within ~50 pc, but none has yet been found.**



Hooper & TL (1803.08046)



# Discovering Pulsars at TeV Energies

- 5 / 39 sources in the 2HWC catalog are correlated with bright, middle-aged (100 — 400 kyr) pulsars.


2HWC Name	ATNF Name	Distance (kpc)	Angular Separation	Projected Separation	Expected Flux ( $\times 10^{-15}$ )	Actual Flux ( $\times 10^{-15}$ )	Flux Ratio	Expected Extension	Actual Extension	Age (kyr)	Chance Overlap
J0700+143	B0656+14	0.29	0.18°	0.91 pc	43.0	23.0	1.87	2.0°	1.73°	111	0.0
J0631+169	J0633+1746	0.25	0.89°	3.88 pc	48.7	48.7	1.0	2.0°	2.0°	342	0.0
J1912+099	J1913+1011	4.61	0.34°	27.36 pc	13.0	36.6	0.36	0.11°	0.7°	169	0.30
J2031+415	J2032+4127	1.70	0.11°	3.26 pc	5.59	61.6	0.091	0.29°	0.7°	181	0.002
J1831-098	J1831-0952	3.68	0.04°	2.57 pc	7.70	95.8	0.080	0.14°	0.9°	128	0.006

- 12 others with young pulsars
  - 2.3 chance overlaps
  - TeV emission may be contaminated by SNR

2HWC Name	ATNF Name	Distance (kpc)	Angular Separation	Projected Separation	Expected Flux ( $\times 10^{-15}$ )	Actual Flux ( $\times 10^{-15}$ )	Flux Ratio	Expected Extension	Actual Extension	Age (kyr)	Chance Overlap
J1930+188	J1930+1852	7.0	0.03°	3.67 pc	23.2	9.8	2.37	0.07°	0.0°	2.89	0.002
J1814-173	J1813-1749	4.7	0.54°	44.30 pc	243	152	1.60	0.11°	1.0°	5.6	0.61
J2019+367	J2021+3651	1.8	0.27°	8.48 pc	99.8	58.2	1.71	0.28°	0.7°	17.2	0.04
J1928+177	J1928+1746	4.34	0.03°	2.27 pc	8.08	10.0	0.81	0.11°	0.0°	82.6	0.002
J1908+063	J1907+0602	2.58	0.36°	16.21 pc	40.0	85.0	0.47	0.2°	0.8°	19.5	0.26
J2020+403	J2021+4026	2.15	0.18°	6.75 pc	2.48	18.5	0.134	0.23°	0.0°	77	0.01
J1857+027	J1856+0245	6.32	0.12°	13.24 pc	11.0	97.0	0.11	0.08°	0.9°	20.6	0.06
J1825-134	J1826-1334	3.61	0.20°	12.66 pc	20.5	249	0.082	0.14°	0.9°	21.4	0.14
J1837-065	J1838-0655	6.60	0.38°	43.77 pc	12.0	341	0.035	0.08°	2.0°	22.7	0.48
J1837-065	J1837-0604	4.78	0.50°	41.71 pc	8.3	341	0.024	0.10°	2.0°	33.8	0.68
J2006+341	J2004+3429	10.8	0.42°	80.07 pc	0.48	24.5	0.019	0.04°	0.9°	18.5	0.08




# Emission Morphologies





# Evolutionary History of Millisecond Pulsars






# Dark Matter Thermalization

Bertoni et al. (2013; 1309.1721)

- **Dark Matter thermalization is always suppressed by Pauli blocking.**
- **Superfluidity and superconductivity effects in the NS core also have a sizable effect.**
- **However, if DM is trapped within the NS, interactions are inevitable. in pessimistic scenarios, DM thermalizes in a timeframe:**



$$t_{th} \simeq 3.7 \text{ kyr} \frac{\frac{m_X}{m_B}}{(1 + \frac{m_X}{m_B})^2} \left( \frac{2 \times 10^{-45} \text{ cm}^2}{\sigma_{nX}} \right) \left( \frac{10^5 \text{ K}}{T_{NS}} \right)^2$$



# Dark Matter Collapse

---

- ▶ Two paths are possible:
  - ▶ **If dark matter can annihilate**, the large densities make annihilation inevitable.
  - ▶ **If dark matter cannot annihilate**, dark matter builds mass until it exceeds its own degeneracy pressure. For Fermionic dark matter this is:

$$M_{crit}^{ferm} \simeq M_{pl}^3 / m_X^2$$


- ▶ It then collapses on a timescale:

$$\begin{aligned} \tau_{co} &\simeq \frac{1}{n\sigma_{nx}v_x} \left( \frac{p_F}{\Delta p} \right) \left( \frac{m_x}{2m_n} \right) \\ &\simeq 4 \times 10^5 \text{ yrs} \left( \frac{10^{-45} \text{ cm}^2}{\sigma_{nx}} \right) \left( \frac{r_x}{r_0} \right), \end{aligned}$$



# The Missing Pulsar Problem

Dexter, O'Leary (1310.7022)



- ▶ Large pulse dispersion was reasonable culprit

$$\Delta\tau \sim 1 \text{ s} \left( \frac{\text{Ghz}}{\nu} \right)^4$$

- ▶ Magnetar found in X-Ray observations in 2013.

- ▶ No pulse dispersion in X-Rays

- ▶ Magnetar subsequently found in radio
- ▶ Pulse dispersion is small!
- ▶ Why aren't any other pulsars observed !?



# Gravitational Waves from NS Collapse


## ► Gravitational Waves from DM induced collapse

$$h_c \sim 5 \times 10^{-22} \left( \frac{M}{M_\odot} \right) \left( \frac{10 \text{ kpc}}{D} \right) @ 531 \text{ Hz},$$

Baiotti et al. (gr-qc/0701043)

## ► Single NS collapse models have been considered (primarily from accretion induced collapse).



## ► DM induced NS collapse observable throughout the Milky Way ( $0.01 \text{ yr}^{-1}$ ?)





# Differentiating Black Hole and NS Mergers



- ▶ **Anomalies in the tidal strain of binary neutron star mergers.**
  - ▶ **DM induced NS collapse produces a population of 1.4  $M_{\odot}$  black holes.**
  - ▶ **Can potentially see differences in merger and ring-down, but not presently feasible.**






# Particle Physics Mash-Up

M A S C U P




- Low mass axions can mediate forces between inspiraling neutron stars, providing effects comparable to gravity.
- LIGO observations can probe the low-mass axion window.





	Description	Symbol	Value
Independent parameters	Physical baryon density parameter <sup>[a]</sup>	$\Omega_b h^2$	$0.022\,30 \pm 0.000\,14$
	Physical dark matter density parameter <sup>[a]</sup>	$\Omega_c h^2$	$0.1188 \pm 0.0010$
	Age of the universe	$t_0$	$13.799 \pm 0.021 \times 10^9$ years
	Scalar spectral index	$n_s$	$0.9667 \pm 0.0040$
	Curvature fluctuation amplitude, $k_0 = 0.002 \text{ Mpc}^{-1}$	$\Delta_R^2$	$2.441^{+0.088}_{-0.092} \times 10^{-9}$ <sup>[17]</sup>
	Reionization optical depth	$\tau$	$0.066 \pm 0.012$







parameters	Scalar spectral index	$n_s$	$0.9667 \pm 0.0040$
	Curvature fluctuation amplitude, $k_0 = 0.002 \text{ Mpc}^{-1}$	$\Delta_R^2$	$2.441^{+0.088}_{-0.092} \times 10^{-9}$ <a href="#">[17]</a>
	<a href="#">Reionization optical depth</a>	$\tau$	$0.066 \pm 0.012$







meters	Curvature fluctuation amplitude, $k_0 = 0.002 \text{ Mpc}^{-1}$	$\Delta_R^2$	$2.441^{+0.088}_{-0.092} \times 10^{-9}$ <sup>[17]</sup>
	Reionization optical depth	$\tau$	$0.066 \pm 0.012$








# One Slide On WIMP Dark Matter



- The standard WIMP freeze-out scenario is still the best-motivated model to explain the Dark Matter abundance.




# Direct Detection






# Dark Matter Flux Depends on NS EOS





# Axion and Neutrino Cooling in Neutron Stars




Hamaguchi et al. (1806.07151)




# r-process Enrichment

Margutti et al. (1801.03531)






# What is the r-process?




## The Origin of the Solar System Elements




Graphic created by Jennifer Johnson

Astronomical Image Credits:  
ESA/NASA/AASNova





# Fast Radio Bursts





# One More Slide on Axion Dark Matter

Axions proposed to solve the strong-CP problem


$$L_\theta = \theta \frac{g^2}{32\pi^2} F_a^{\mu\nu} \tilde{F}_{a\mu\nu}$$

This is the sum of two different terms, that independently must be small.

$$|\Theta_{\text{QCD}} + \arg \det M_q| \lesssim 10^{-9}$$

This provides you with an independent way to solve the strong-CP problem - by setting  $m_u = 0$ .

However, this appears to be at odds with experimental data.





# QCD Axion Mass Bounds

---

**QCD Axion obtains its mass from its decay constant and the coupling to quarks:**

$$m_a = \frac{f_\pi m_\pi}{f_a} \left( \frac{z}{(1+z+w)(1+z)} \right)^{1/2} \\ = 0.60 \text{ eV} \frac{10^7 \text{ GeV}}{f_a},$$

**The high-mass range of the QCD axion (low- $f_a$ ) is set by the constraint that the axion never comes into thermal equilibrium (and light-through a wall and astrophysical constraints).**


**The low mass (high- $f_a$ ) limit is set such that axions don't overclose the universe:**

$$\Omega_a h^2 \approx 0.23 \times 10^{\pm 0.6} (f_a / 10^{12} \text{ GeV})^{1.175} \Theta_i^2 F(\Theta_i)$$



# Goldreich-Julian Current

Kalapothisarakos et al. (1108.2138)



NS is a conductor with a strong rotating magnetic field. Thus, an electric field and current are formed.

In these simulations the NS is not a perfect dipole.



# A Signal

10% of Star Formation in central 200 pc of Milky Way

## Massive Star Formation in the Galactic Center

By Don F. Figer

Rochester Institute of Technology, Rochester, NY, USA

The Galactic center is a hotbed of star formation activity, containing the most massive star formation site and three of the most massive young star clusters in the Galaxy. Given such a rich environment, it contains more stars with initial masses above  $100 M_{\odot}$  than anywhere else in the Galaxy. This review concerns the young stellar population in the three massive stellar clusters, the population of younger stars in the field population, as it relates to massive star formation above  $100 M_{\odot}$  in the Galactic center. The stars surrounding the central black hole, and the bulk of the stars in the field population. The fossil record in the Galactic center suggests that the recently formed massive stars represent-day examples of similar populations that must have been formed through episodes stretching back to the time period when the Galaxy was forming.

### Introduction

Galactic center (GC) is an important region in many models. It contains a tiny fraction of the Galaxy's mass, but a large fraction of its star formation activity.



# Dark Matter Induced Collapse in dSphs

Bramante & TL (1601.06784)

- **The dispersion velocity in dwarfs is also small.**
  - **Reticulum II:  $3.3 \pm 0.7 \text{ km s}^{-1}$  (Simon et al. 2015)**
- **Dark matter accumulation rate scales inversely with velocity:**


$$\dot{m}_x = \pi \rho_x \frac{2GM R}{v_x} \left(1 - \frac{2GM}{R}\right)^{-1}$$
$$\simeq \frac{10^{26} \text{ GeV}}{\text{s}} \left(\frac{\rho_x}{\text{GeV/cm}^3}\right) \left(\frac{200 \text{ km/s}}{v_x}\right),$$

- **Dwarf Spheroidal Galaxies are an optimal laboratory for asymmetric dark matter detection.**



# Dark Matter Induced Collapse in dSphs



- **Reticulum II dSph**
  - **Discovered by DES in 2015**
  - **Spectroscopic follow-up determined r-process abundances.**
  - **Large r-process abundance, but low metallicity!**
- **Points to a rare formation channel (NS mergers)**





# Dark Matter Induced Collapse in dSphs

- **Normalize the nuclear cross-section to the missing pulsar problem.**
- **Supernovae produce ~100 events.**
- **Mergers produce ~0.0005 events**
- **DM induced collapse produces ~0.1-3 events.**






# Dark Matter Induced Collapse in dSphs

Roederer 2011 (1104.5056)

- **Prediction: Globular Clusters should not be similarly r-process enriched.**
- **In fact, no globular cluster has been observed to have an r-process overabundance exceeding 1.2 dex.**
- **6 of 9 stars in Reticulum II have r-process enrichment exceeding 1.68 dex.**





# Dark Matter Induced Collapse in dSphs

position dependent

$$M_{DM} \propto \sigma_{xp}$$

$t_{NS}$

$$\left( \frac{\rho_x}{v_x} \right)$$


particle physics

in any  
p.p. model

$$t_{imp}(r) \propto \frac{\rho_x}{v_x}$$




# Pulsars Produce the Positron Excess



- Can calculate the gamma-ray spectrum necessary to fit the Geminga data from HAWC and Milagro
- Can use this to calculate the underlying steady-state electron and positron spectrum



# Pulsars Produce the Positron Excess





- Utilizing a diffusion model, along with the steady state electron spectrum, and the morphology of the emission, can calculate the fraction of the electron energy lost before escaping the halo.
- **Less energetic electrons make it to the ISM!**



# Pulsars Produce the Positron Excess

- In these models, Geminga naturally produces ~50% of the positron excess.
- The total contribution from the remaining Milky Way pulsars produces the remaining emission.
- Difficult to understand TeV halo spectrum if pulsars do not make the positron excess.






# Pulsars Produce the TeV Excess

- **Milagro detected bright diffuse TeV emission along the Galactic plane.**

- **The intensity of this emission is incompatible with hadronic models constrained by Fermi and Argo-YBJ data.**





TL & Buckman (1707.01905)

- **TeV halos produce a hard spectrum component that naturally explains the intensity and spectrum of this emission.**



# Pulsars Produce Anisotropic Diffusion

- **Diffusion near TeV halo is known to be suppressed - by two orders of magnitude!**
- **Diffusion constant near us must be high to explain observations of 10 TeV electrons.**
- **Pulsars produce regions of low-diffusion, where TeV halos shine!**






# Pulsars Produce Anisotropic Diffusion

- **Cosmic-Ray electrons produced by the pulsar obtain a steep gradient.**
- **This excites Alfven waves moving parallel to the electron gradient.**



$$\Gamma_{\text{CR}}(k) = \frac{2\pi}{3} \frac{c|v_A|}{k\mathcal{W}(k)U_0} \left[ p^4 \frac{\partial f}{\partial z} \right]_{p_{\text{res}}}$$



- **These Alfven waves dominate cosmic-ray turbulence, because they are resonant with the electron energy - leads to low diffusion**



# Pulsars Produce Galactic Center Pevatron



- **The spectrum of the HESS pevatron looks like the Geminga spectrum.**
- **Diffuse electrons can be made via pulsar natal kicks.**



# One Slide on Axion Dark Matter

Axions proposed to solve the strong-CP problem

$$L_\theta = \theta \frac{g^2}{32\pi^2} F_a^{\mu\nu} \tilde{F}_{a\mu\nu}$$

If this constant is promoted to a field, its self-interactions drive it to 0:

$$\mathcal{L}_\Theta \rightarrow \mathcal{L}_a = \frac{1}{2} (\partial_\mu a)^2 - \frac{\alpha_s}{8\pi f_a} a G \tilde{G}$$

This term must couple to the EM field, allowing for decays to photons:

$$\mathcal{L}_{\text{int}} = -\frac{1}{4} g_{a\gamma} F_{\mu\nu} \tilde{F}^{\mu\nu} a = g_{a\gamma} \mathbf{E} \cdot \mathbf{B} a$$

